



Reduced field efficacy of ivermectin against *Ostertagia ostertagi* and *Cooperia oncophora* in Danish cattle

Peña-Espinoza, Miguel; Drag, Markus; Hansen, Tina Vicky Alstrup; Thamsborg, Stig Milan; Enemark, Heidi Larsen

Publication date:
2015

Document version
Early version, also known as pre-print

Citation for published version (APA):
Peña-Espinoza, M., Drag, M., Hansen, T. V. A., Thamsborg, S. M., & Enemark, H. L. (2015). *Reduced field efficacy of ivermectin against Ostertagia ostertagi and Cooperia oncophora in Danish cattle*. Abstract from 25th International Conference of the World Association for the Advancement of Veterinary Parasitology, Liverpool, United Kingdom. http://orbit.dtu.dk/files/117697125/Pe_a_Espinoza_et_al_FECRT_WAAVP2015.pdf

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/282976160>

Reduced field efficacy of ivermectin against *Ostertagia ostertagi* and *Cooperia oncophora* in Danish cattle

Conference Paper · August 2015

READS

23

5 authors, including:



Miguel Peña-Espinoza

Technical University of Denmark

9 PUBLICATIONS 4 CITATIONS

SEE PROFILE



Tina Hansen

University of Copenhagen

13 PUBLICATIONS 48 CITATIONS

SEE PROFILE



Stig Milan Thamsborg

University of Copenhagen

193 PUBLICATIONS 3,863 CITATIONS

SEE PROFILE



Heidi Larsen Enemark

Norwegian Veterinary Institute

59 PUBLICATIONS 909 CITATIONS

SEE PROFILE

Thursday 20 August 2015

Concurrent Session 9.1

A: Drugs and drug resistance

Ref: O225/0639

Reduced field efficacy of ivermectin against *Ostertagia ostertagi* and *Cooperia oncophora* in Danish cattle

Miguel Peña-Espinoza^{1,2}, Markus Drag¹, Tina Hansen², Stig Thamsborg², Heidi Enemark^{1,3}

¹National Veterinary Institute, Technical University of Denmark, Copenhagen, Denmark,

²Department of Veterinary Disease Biology, University of Copenhagen, Copenhagen, Denmark,

³Norwegian Veterinary Institute, Oslo, Norway

Field reports of anthelmintic resistance against the widely-used macrocyclic lactones (ML) in gastrointestinal nematodes (GIN) of cattle have appeared in NW-Europe in recent years.

Objective: The aim of this study was to assess the efficacy of ivermectin (IVM) against field infections with GIN in Danish cattle. In addition, we evaluated a novel quantitative (q) real-time PCR assay for accurate identification of surviving nematode species after treatment.

Methods: Six farms were selected based on mean faecal egg counts (FEC) of ≥ 100 nematode eggs per gram in first-season grazing heifers. All selected farms had a history of use of ML. In each farm, 20 heifers were selected for faecal egg count reduction test (FECRT) and individual weights were measured using a girth-tape. Ten animals were treated (s.c.) with 0.2 mg IVM/kg body weight and ten animals were left untreated as controls. FEC were investigated in all animals at the day of treatment (0) and at days 14 and 21 post-treatment (p.t). In parallel, L3 were cultured from pooled faeces from all animals in each group. DNA isolated from pooled L3 was analysed by a qPCR quantifying copies of the second internal transcribed spacer (ITS2) specific for *Ostertagia ostertagi* and *Cooperia oncophora*. FEC reduction percentages (FECR%) in IVM-treated groups were calculated following WAAVP guidelines including FECs of the control group. Reduced efficacy was defined when $\text{FECR}\% \leq 95\%$ and lower confidence interval [CI] $\leq 90\%$.

Results: At day 14 p.t., FECR% of IVM treatments in all six farms varied from 75% to 96% (lower CI ranging from 39% to 91%). At day 21 p.t., IVM treatments showed a reduced efficacy in all farms, with FECR% varying from 45% to 91% (lower CI ranging from -34% to 65%). At day 0, ITS2 copies from *O. ostertagi* and *C. oncophora* were identified in larval cultures from treated groups in all six farms. At day 14 p.t. in treated animals, *O. ostertagi* L3 were found in 1 farm and *C. oncophora* L3 in 3 farms. On day 21 p.t., *O. ostertagi* L3 and *C. oncophora* L3 were detected in treated animals of 2 and 4 farms, respectively.

Discussion: Reduced field efficacy of IVM was confirmed by FECRT in 5 out of 6 farms at day 14 p.t. and in all farms at day 21 p.t. The qPCR was able to identify *O. ostertagi* and *C. oncophora* populations surviving IVM treatment. This emphasizes the need of monitoring the efficacy of anthelmintic treatments in cattle using sensitive methods.